

Fast implementation of the Tukey depth *

Xiaohui Liu^a

^a School of Statistics, Jiangxi University of Finance and Economics, Nanchang, Jiangxi 330013, China

Abstract. This paper presents two new algorithms for fast calculating the Tukey depth of dimensions $p \geq 3$. The first algorithm is naive, and can compute *exactly* the Tukey depth of a single point with complexity $O(n^{p-1} \log(n))$. Compared to the first, the second algorithm further utilizes the *quasiconcave* of the Tukey depth function, and hence can be implemented with complexity less than $O(\frac{1}{2^{p-2}} n^{p-1} \log(n))$. Both of them are of *combinatorial* properties. Experiments show that the proposed algorithms require quite minimum memory and run much faster than the existing procedures.

Key words: Tukey depth; Quasiconcave; Combinatorial property; Fast computation

2000 Mathematics Subject Classification Codes: 62F10; 62F40; 62F35

1 Introduction

To provide a desirable ordering for multivariate data, [Tukey \(1975\)](#) heuristically proposed a useful tool of the statistical depth function. With respect to the distribution P of X in $\mathcal{R}^p$ ($p \geq 1$), he defined the Tukey depth of a point z as the minimum probability mass carried by any closed halfspace containing z . Usually, the Tukey depth function is expressed as

$$D(z|P) = \inf_{u \in S^{p-1}} P(u^\top X \leq u^\top z),$$

where $S^{p-1} = \{v \in \mathcal{R}^p : \|v\| = 1\}$. For n given observations $\mathcal{X}^n := \{X_i\}_{i=1}^n$ in $\mathcal{R}^p$, its sample version is correspondingly

$$D_{np}(z) := D(z|P_n) = \inf_{u \in S^{p-1}} P_n(u^\top X \leq u^\top z), \quad (1)$$

where P_n denotes the empirical distribution of $\mathcal{X}^n$.

The Tukey depth has proved very desirable. It satisfies all four properties that define a general notion of statistical depth functions, namely, affine invariance, maximality at center, monotonicity relative to deepest point, and vanishing at infinity ([Zuo and Serfling,](#)

*Corresponding author's email: csulixh912@gmail.com

2000). In practice, it finds many applications in cases such as confidence region constructions (Yeh and Singh, 1997) and classifications (Li *et al.*, 2012). It even can characterize the underlying distribution under very mild conditions (Kong and Zuo, 2010). Latest developments indicate that the Tukey depth also has a very strong connection with the multiple-output quantile regression methodology (Hallin *et al.*, 2010; Kong and Mizera, 2012).

However, its *exact* computation is challenging. This is mainly because the function $P_n(u^\top X \leq u^\top z)$ involved is discontinuous and non-convex with respect to $u \in \mathcal{S}^{p-1}$. Hence, it is difficult to utilize the conventional optimization methods to obtain the infimum of $P_n(u^\top X \leq u^\top z)$. To make the computation possible, special attention should be paid to how to reduce the number of u . The first result in that direction is made by Rousseeuw and Ruts (1996) for *bivariate data* relying on the idea of the circular sequence (Edelsbrunner, 1987). Using a similar idea, Rousseeuw and Struyf (1998) constructed an exact algorithm for computing the Tukey depth for $p = 3$.

For *more high dimensional data*, Liu and Zuo (2014a) developed a feasible cone enumeration procedure based on the breadth-first search algorithm. The cones considered in Liu and Zuo (2014a) satisfy that their vertexes contain all $\binom{n}{p-1}$ *critical direction vectors*, which are normal to the hyperplanes passing through $\{z, X_{i_1}, \dots, X_{i_{p-1}}\}$, where $i_1, \dots, i_{p-1}$ are distinct and $i_1, \dots, i_{p-1} \in \{1, 2, \dots, n\}$. Recently, Mozharovskiy (2014) further refined the algorithm of Liu and Zuo (2014a). They found that it is possible to compute the Tukey depth by directly considering these $\binom{n}{p-1}$ *critical direction vectors*. Since Dyckerhoff and Mozharovskiy's approach is of combinatorial nature, and does not need to take into account any space ordering, the implementation requires much less memory and runs much faster.

In this paper, we further improve Dyckerhoff and Mozharovskiy's procedure. We find that it is useful to extend the definition of the Tukey depth for a single point z into the version for a subspace $\mathbf{V}$. Then relying on this, we propose for dimensions $p \geq 3$ a new exact algorithm which is still of combinatorial nature, but possesses exactly the complexity of $O(n^{p-1} \log n)$, better than that $O(n^p)$ of Mozharovskiy (2014).

Nevertheless, like all algorithms mentioned above, this new algorithm still needs to fully address all $\binom{n}{p-1}$ *critical direction vectors*. On the other hand, when computing the Tukey depth, we are in fact searching for the infimum of $P_n(u^\top X \leq u^\top z)$ with respect to u . A great proportion of the *critical direction vectors* may be *redundant* in the sense that some of them have values $P_n(u^\top X \leq u^\top z)$ larger than τ , which we assume to be an approximation of the Tukey depth obtained through the approximate methods. A nature question here is whether or not we can eliminate some of them from consideration.

The answer is positive. With the extended definition above, we find it is possible to

utilize the *quasiconcave*, i.e., all depth regions are convex (Mosler, 2013), of the Tukey depth function to reduce greatly the number of *critical direction vectors* involved. An *iterative* algorithm is constructed to realize this idea. This approach is still of combinatorial property because it is been strictly limited to consider direction vectors among the *critical direction vectors*. Hence it can be implemented very efficiently. This algorithm is depth-depending. The smaller the depth value of the point z is, the less the computation time this algorithm tends to consume.

The proposed algorithms have been implemented in Matlab. The whole code can be obtained through emailing to the author. Data examples are also provided to illustrate the performance of the proposed algorithms.

The rest of this paper is organized as follows. Section 2 extends the conventional definition of the Tukey depth for a single point z to the version for a subspace. Section 3 provides a refined *combinatorial* algorithm for *exactly* computing the Tukey depth. Section 4 develops an *adaptively iterative* procedure. Several data examples are given in Section 5 to illustrate the performance of the proposed algorithms. Section 6 ends the whole paper with some concluding discussions.

2 Tukey depth for a subspace

In the literature, it is known that constructing algorithms directly based on (1) is difficult to utilize some information, such as *quasiconcave*, of the Tukey depth function to fasten the computation. To this end, we propose to consider a useful extended version of (1) in the following.

Without loss of generality, we assume that X^p are in general position (Mosler et al., 2009). Note that $D_{np}(z|P_n) = D_{np}(0|P_{nz})$ for any given $z \in \mathcal{R}^p$ by the affine invariance, in the sequel we suppose that $z = 0$, and pretend the real observations to be $\mathbf{X}^n := \{\mathbf{x}_i\}_{i=1}^n$ with the empirical distribution being P_{nz} , where $\mathbf{x}_i = X_i - z$.

Let $\mathbf{W}^\perp$ be the orthogonal complement of the subspace $\mathbf{W}$. Then for a r -dimensional subspace $\mathbf{V}_r$ of $\mathcal{R}^p$ ($0 \leq r < p$), we define its Tukey depth value with respect to $\mathbf{X}^n$ as follows:

$$D_{np}(\mathbf{V}_r) = \inf_{u \in \mathcal{S}^{p-1} \cap \mathbf{V}_r^\perp} P_n^\perp(u^\top X_r^\perp \leq 0), \quad (2)$$

where $X_r^\perp$ denotes the projection of X onto the $(p - r)$ -dimensional subspace $\mathbf{V}_r^\perp$, and $P_n^\perp$ the empirical distribution of $\{\mathbf{x}_{ri}^\perp | \mathbf{x}_{ri}^\perp \text{ the projection of } \mathbf{x}_i, \mathbf{x}_i \in \mathbf{X}^n\}$. When $r = 0$, we assume that $\mathbf{V}_r$ contains only a single point $\{0\}$, and its orthogonal complement subspace is the whole $\mathcal{R}^p$. In this sense, $D_{np}(\mathbf{V}_r)$ can be regarded as an extension of (1).

Clearly, for a given $\mathbf{V}_r$ ($0 < r < p$),

$$D_{np}(\mathbf{V}_r) = \inf_{u \in \mathcal{S}^{p-1} \cap \mathbf{V}_r^\perp} P_n^\perp(u^\top X_r^\perp \leq 0) = \inf_{u \in \mathcal{S}^{p-1} \cap \mathbf{V}_r^\perp} P_n(u^\top X \leq 0) \geq D_{np}(0).$$

Based on this, when $\mathbf{V}_r$ is any given, it is trivially to show that

$$D_{np}(0) = \inf_{\mathbf{V}_r \in \mathcal{V}_r} D_{np}(\mathbf{V}_r), \quad (3)$$

where $\mathcal{V}_r$ denotes the set containing all r -dimensional subspaces. When $r = p - 1$, (3) deduces to

$$D_{np}(0) = \inf_{u \in \mathcal{S}^{p-1}} D_{np}(\mathbf{H}_u)$$

with $\mathbf{H}_u = \{x \in \mathcal{R}^p \mid u^\top x = 0, u \in \mathcal{S}^{p-1}\}$.

For $p \geq 2$, [Mozharovskiy \(2014\)](#) have recently showed that $\binom{n}{p-1}$ *critical directors vectors*, normal to the hyperplanes passing through $\{\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{p-1}}, 0\}$, are sufficient to determine the value of $D_{np}(0)$, where $i_1, i_2, \dots, i_{p-1}$ are distinct and $i_1, i_2, \dots, i_{p-1} \in \{1, 2, \dots, n\}$; see also [Liu and Zuo \(2014a\)](#). This result is useful because it transforms the computation of the Tukey depth into an optimal issue over a finite number of *critical direction vectors*.

Equivalently, from the point of view of subspaces, one may conclude that $\binom{n}{p-1}$ subspaces spanned by $\{\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{p-1}}, 0\}$ are sufficient to determine the value of $D_{np}(0)$. In fact, one can reexpress the results of Algorithm 3 and Corollary 8 of [Mozharovskiy \(2014\)](#) as follows:

$$D_{np}(0) = \min_{\mathbf{I}_{p-1} \in \mathcal{I}_{p-1}} D_{np}(\mathbf{S}_{\mathbf{I}_{p-1}}) - \frac{p-1}{n}, \quad (4)$$

which is a special case of the following useful proposition.

Proposition 1. *Assume that the observations $\mathbf{X}^n$ are in general position. Then for any $r = 1, 2, \dots, p-1$, we have that*

$$D_{np}(0) = \min_{\mathbf{I}_r \in \mathcal{I}_r} D_{np}(\mathbf{S}_{\mathbf{I}_r}) - \frac{r}{n},$$

where

$$\mathcal{I}_r = \{\{i_1, i_2, \dots, i_r\} \mid i_1, i_2, \dots, i_r \text{ distinct, and } i_1, i_2, \dots, i_r \in \{1, 2, \dots, n\}\}, \quad (5)$$

and $\mathbf{S}_{\mathbf{I}_r} := \mathbf{S}_{i_1, i_2, \dots, i_r} = \text{span}(\mathbf{x}_l : l = i_1, i_2, \dots, i_r)$ denotes the r -dimensional subspace of $\mathcal{R}^p$ spanned by $\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_r}$.

Proof. By definition, for any $\mathbf{I}_r := \{i_1, i_2, \dots, i_r\} \in \mathcal{I}_r$, we have

$$D_{np}(\mathbf{S}_{\mathbf{I}_r}) = \inf_{u \in \mathcal{S}^{p-1} \cap \mathbf{S}_{\mathbf{I}_r}^\perp} P_n^\perp(u^\top X_r^\perp \leq 0) =: D_{n(p-r), \mathbf{I}_r}(0).$$

For $D_{n(p-r), \mathbf{I}_r}(0)$, similar to (4), we obtain

$$D_{n(p-r), \mathbf{I}_r}(0) = \min_{\mathbf{J}_{p-r-1} \in \mathcal{J}_{p-r-1}} D_{n(p-r), \mathbf{I}_r}(\mathbf{S}_{\mathbf{J}_{p-r-1}}^\perp) - \frac{(p-r)-1}{n},$$

where

$$\begin{aligned} \mathcal{J}_{p-r-1} &= \{\{j_1, j_2, \dots, j_{p-r-1}\} \mid j_1, j_2, \dots, j_{p-r-1} \text{ distinct, and} \\ &\quad j_1, j_2, \dots, j_{p-r-1} \in \{1, 2, \dots, n\} \setminus \{i_1, i_2, \dots, i_r\}\}, \end{aligned}$$

and $\mathbf{S}_{\mathbf{J}_{p-r-1}}^\perp = \text{span}(\mathbf{x}_l^\perp : l = j_1, j_2, \dots, j_{p-r-1})$ denotes the subspace spanned by $\mathbf{x}_{j_1}^\perp, \mathbf{x}_{j_2}^\perp, \dots, \mathbf{x}_{j_{p-r-1}}^\perp$.

Based on this, it is easy to show that

$$\begin{aligned} \min_{\mathbf{I}_r \in \mathcal{I}_r} D_{np}(\mathbf{S}_{\mathbf{I}_r}) &= \min_{\mathbf{I}_r \in \mathcal{I}_r} \left\{ \min_{\mathbf{J}_{p-r-1} \in \mathcal{J}_{p-r-1}} D_{n(p-r), \mathbf{I}_r}(\mathbf{S}_{\mathbf{J}_{p-r-1}}^\perp) - \frac{p-r-1}{n} \right\} \\ &= \min_{\mathbf{I}_{p-1} \in \mathcal{I}_{p-1}} D_{np}(\mathbf{S}_{\mathbf{I}_{p-1}}) - \frac{p-1}{n} + \frac{r}{n} \\ &= D_{np}(0) + \frac{r}{n}. \end{aligned}$$

Then this proposition follows immediately.

Proposition 1 coincides with the result obtained by [Liu and Zuo \(2014a\)](#). That is, for any $u_0 \in \mathcal{S}^{p-1}$ such that

$$D_{np}(0) = \inf_{u \in \mathcal{S}^{p-1}} P_n(u^\top X \leq 0) = P_n(u_0^\top X \leq 0),$$

the hyperplane $\{x \in \mathcal{R}^p \mid u_0^\top x = 0\}$ contains no observation of $\mathcal{X}^n$, if $\mathcal{X}^n \cap \{0\}$ are in general position.

3 A refined combinatorial algorithm

Since the computation of the Tukey depth is trivial when $p = 1$, we focus only on the cases of $p \geq 2$ in the following.

For $p \geq 2$, [Mozharovskiy \(2014\)](#) recently proposed a combinatorial algorithm, whose implementation runs faster and requires much less memory than that constructed on the breadth-first search algorithm. It turns out that their procedure is of the complexity $O(n^p)$. When $p = 2, 3$, the order of the complexity of their procedure is higher than that of few existing algorithms; see for example [Rousseeuw and Ruts \(1996\)](#) and [Rousseeuw and Struyf \(1998\)](#).

If carefully investigating Dyckerhoff and Mozharovskyi's algorithm, it is easy to find that their proposal computes $D_{np}(0)$ actually through finding the minimum:

$$\min_{\mathbf{I}_{p-1} \in \mathcal{I}_{p-1}} D_{np}(\mathbf{S}_{\mathbf{I}_{p-1}}) - \frac{p-1}{n}$$

over all $(p-1)$ -dimensional subspace $\mathbf{S}_{\mathbf{I}_{p-1}}$ spanned by $\binom{n}{p-1}$ combinations $\{\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{p-1}}, 0\}$. On the other hand, Proposition 1 indicates that we can also utilize the fact that $D_{np}(0) = \min_{\mathbf{I}_r \in \mathcal{I}_r} D_{np}(\mathbf{S}_{\mathbf{I}_r}) - \frac{r}{n}$ to compute $D_{np}(0)$ with an another $r (\neq p-1)$.

Among $r = 1, 2, \dots, p-2$, our favourite is

$$D_{np}(0) = \min_{\mathbf{I}_{p-2} \in \mathcal{I}_{p-2}} D_{np}(\mathbf{S}_{\mathbf{I}_{p-2}}) - \frac{p-2}{n}.$$

The reasons are as follows. There are only $\binom{n}{p-2}$ combinations $\mathbf{I}_{p-2}$. For each $\mathbf{I}_{p-2}$, $D_{np}(\mathbf{S}_{\mathbf{I}_{p-2}}) = D_{n2, \mathbf{I}_{p-2}}(0)$, which is in fact a bivariate Tukey depth. While for bivariate data, it is known that the complexity of some well-developed algorithms is only of the order $O(n \log(n))$ (Rousseeuw and Ruts, 1996). Hence, Dyckerhoff and Mozharovskyi's algorithm may be further improved to the version with complexity $O(n^{p-1} \log(n))$.

The detailed procedure is described as follows. It is actually a further refined version of the combinatorial algorithm suggested by Mozharovskyi (2014).

Algorithm 1. (for p -dimensional data with $p \geq 3$)

Input: $\mathbf{X}^n = \{\mathbf{x}_1, \dots, \mathbf{x}_n\} \in \mathcal{R}^p$, $3 \leq p < n < \infty$, $\mathbf{X}^n$ in general position.

Step 1. Let $N_{\min} = n$. For each $\mathbf{I}_{p-2} := \{i_1, \dots, i_{p-2}\} \in \mathcal{I}_{p-2}$ (see (5)), do:

- (a) compute two orthogonal bases e_1 and e_2 of the orthogonal complement subspace of that spanned by $\{0, \mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{p-2}}\}$,
- (b) compute the bivariate Tukey depth $D_{n2}(0)$ with respect to $\{\mathbf{y}_i := (e_1^\top \mathbf{x}_i, e_2^\top \mathbf{x}_i)^\top, i = 1, 2, \dots, n\}$,
- (c) if $N_{\min} > nD_{n2}(0)$, then $N_{\min} = nD_{n2}(0)$.

Step 2. Return $\frac{N_{\min} - (p-2)}{n}$.

Output: $\frac{N_{\min} - (p-2)}{n}$.

Note that the computation of the bivariate Tukey depth $D_{n2}(0)$ is very key for Algorithm 1. A good procedure of computing the bivariate Tukey depth $D_{n2}(0)$ can make the whole algorithm implemented very fast. Therefore, we propose to consider the following approach, which is very efficient, and has been tested by us in many experiments.

Algorithm 2. (for bivariate data only)

Input: $\mathbf{Y}^n = \{\mathbf{y}_1, \dots, \mathbf{y}_n\} \in \mathcal{R}^2$.

Step 1. Let $M_{\min} = n$. Do:

- (a) compute $m = \#\{i \mid \mathbf{y}_i \neq 0, i = 1, \dots, n\}$, and let $\widetilde{\mathbf{Y}}^m := \{\widetilde{\mathbf{y}}_i\}_{i=1}^m = \{\mathbf{y}_i \mid \mathbf{y}_i \neq 0, \mathbf{y}_i \in \mathbf{Y}^n\}$, where $\#\mathcal{A}$ denotes the cardinality of the set $\mathcal{A}$,
- (b) for each $i \in \{1, 2, \dots, m\}$, compute the polar angles $\theta_i \in [0, 2\pi)$ of the direction vector normal to the line passing through 0 and $\widetilde{\mathbf{y}}_i$, set $\widetilde{\theta}_i = \theta_i$ if $\theta_i \in [0, \pi)$ and $\widetilde{\theta}_i = \theta_i - \pi$ otherwise,
- (c) for $\mathbf{u}_0 = (1, 0)^\top$, compute the dichotomies $(L_1, L_2, \dots, L_m)$ and $S_0 = \min\{S_1, S_2\}$, where $S_1 = \sum_{i=1}^m L_i$, and $S_2 = m - S_1$ with $L_i = 1$ if $\mathbf{u}_0^\top \widetilde{\mathbf{y}}_i \geq 0$, else $L_i = 0$ for $i = 1, \dots, m$,
- (d) if $M_{\min} > S_0$, let $M_{\min} = S_0$,
- (e) compute the permutation $(j_1, j_2, \dots, j_m)$ such that $\widetilde{\theta}_{j_1} \leq \widetilde{\theta}_{j_2} \leq \dots \leq \widetilde{\theta}_{j_m}$, and for each $k = 1 : m$, do:
 - (i) if $L_{j_k} = 0$, set $S_1 = S_1 + 1$, else set $S_1 = S_1 - 1$,
 - (ii) compute $S_2 = m - S_1$, and update $S_0 = \min\{S_1, S_2\}$,
 - (iii) if $M_{\min} > S_0$, let $M_{\min} = S_0$.

Step 2. Return $\frac{M_{\min} + n - m}{n}$.

Output: $\frac{M_{\min} + n - m}{n}$.

This procedure is a refined version of the algorithm developed in [Liu and Zuo \(2014a\)](#) when $p = 2$. It is based on the idea of the circular sequence. Since during the computation we only need to sort once a sequence of length m ($m \leq n$, see Step 1-(e)), its complexity is of the order $O(n \log(n))$. By using the dichotomies $(L_1, L_2, \dots, L_m)$ and the permutation $(j_1, j_2, \dots, j_m)$ obtained through sorting, we can easily compute the Tukey depth. In the experiments, we found that it runs even *faster than* the classical algorithm suggested by [Rousseeuw and Ruts \(1996\)](#). Hence, we recommend to use it in Algorithm 1 to compute the bivariate Tukey depth.

4 An adaptive iterative algorithm

Most of the existing procedures have to fully address $\binom{n}{p-1}$ critical direction vectors, no matter where the point $\mathbf{z}$ is located at. On the other hand, a great proportion of these vectors may be redundant, because when computing the Tukey depth, we are in fact searching for the infimum of $P_n(\mathbf{u}^\top \mathbf{X} \leq \mathbf{u}^\top \mathbf{z})$ with respect to $\mathbf{u}$.

This may easily be seen from the illustration given in Figure 1. In this illustration, we are computing the Tukey depth of 0 with respect to a data set containing 10 observations. Assume that we have known that the depth value of 0 is *at most 0.2 through some approximate methods*. Then it is easy to see that the *critical direction vectors* normal to Lines 1-6 are *redundant*, because using them can not produce a smaller depth value than 0.2. Hence it's better to eliminate them from consideration as many as possible. This idea has been utilized by [Johnson et al. \(1998\)](#) for *bivariate* data.




Figure 1: Shown is an illustration of redundant direction vectors for computing the Tukey depth. Clearly, the critical direction vectors normal to Lines 1-6 are *redundant* if we have known that the depth of 0 is at most 0.2.

In this section, we are interested to present an *iterative* procedure for spaces with dimension $p \geq 3$. The most outstanding of this procedure is that it can *adaptively* eliminate many *redundant critical direction vectors* from consideration based on the former iteration.

Before proceeding further, let's provide two useful propositions as follows.

Proposition 2. For an any subspace $\mathbf{V}_r$ ($0 < r < p$) of $\mathcal{R}^p$ ($p > 1$), we have that

$$\sup_{x \in \mathbf{V}_r} D_{np}(x) \leq D_{np}(\mathbf{V}_r).$$

That is, if $D_{np}(\mathbf{V}_r) < \tau$, then all points contained in $\mathbf{V}_r$ lie outside of the τ -th Tukey depth region $\mathcal{D}_n(\tau) = \{x \in \mathcal{R}^p \mid D_{np}(x) \geq \tau\}$.

Proof. Let $\mathbf{B}_\rho = \{x \mid x \in \mathcal{R}^p, \|x\| \leq \rho\}$. Clearly, $\mathbf{B}_\rho$ is compact. Then, by the fact $D_{np}(x) \rightarrow 0$ as $\|x\| \rightarrow \infty$, we show that, for a large enough ρ_0 , there exists an $x_0 \in \mathbf{V}_r \cap \mathbf{B}_{\rho_0}$ such that

$$\sup_{x \in \mathbf{V}_r} D_{np}(x) = D_{np}(x_0).$$

Similarly, the compactness of $\mathcal{S}^{p-1}$ indicates that there is a $u_0 \in \mathcal{S}^{p-1} \cap \mathbf{V}_r^\perp$ such that

$$D_{np}(\mathbf{V}_r) = P_n^\perp(u_0^\top X_r^\perp \leq 0) = P_n(u_0^\top X \leq u_0^\top x_0) \geq \inf_{u \in \mathcal{S}^{p-1}} P_n(u^\top X \leq u^\top x_0) = D_{np}(x_0).$$

This completes the proof.

Proposition 3. For $p \geq 3$ and an any given combination $(\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{p-2}})$, there is an another observation $\mathbf{x}_{i_{p-1}}$ such that

$$D_{np}(\mathbf{S}_{i_1, i_2, \dots, i_{p-2}}) = P_n(\bar{u}_0^\top X \leq 0) - \frac{1}{n},$$

where $\bar{u}_0$ is normal to the hyperplane passing through $\{0, \mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{p-2}}, \mathbf{x}_{i_{p-1}}\}$. More importantly, for any $j_1, j_2, \dots, j_{p-3}$ distinct and $j_1, j_2, \dots, j_{p-3} \in \{i_1, i_2, \dots, i_{p-2}\}$, it holds

$$D_{np}(\mathbf{S}_{i_1, i_2, \dots, i_{p-2}}) \geq D_{np}(\mathbf{S}_{j_1, j_2, \dots, j_{p-3}, i_{p-1}}).$$

Proof. The first part can be proved trivially by following a similar fashion to that of Propositions 1-2. For the second part, since $\bar{u}_0 \in \mathbf{S}_{i_1, i_2, \dots, i_{p-2}, i_{p-1}}^\perp$, then $\bar{u}_0 \in \mathbf{S}_{j_1, j_2, \dots, j_{p-3}, i_{p-1}}^\perp$ holds for any $j_1, j_2, \dots, j_{p-3} \in \{i_1, i_2, \dots, i_{p-2}\}$. Using this, we obtain

$$\begin{aligned} D_{np}(\mathbf{S}_{i_1, i_2, \dots, i_{p-2}}) &= P_n(\bar{u}_0^\top X \leq 0) - \frac{1}{n} \\ &\geq \inf_{u \in \mathcal{S}^{p-1} \cap \mathbf{S}_{j_1, j_2, \dots, j_{p-3}, i_{p-1}}^\perp} P_n^\perp(u_0^\top X_r^\perp \leq 0) \\ &= D_{np}(\mathbf{S}_{j_1, j_2, \dots, j_{p-3}, i_{p-1}}). \end{aligned}$$

These two propositions, together with the *quasiconcave* of the Tukey depth function, indicate that $\mathbf{S}_{j_1, j_2, \dots, j_{p-3}, i_{p-1}}$ lies more outside than $\mathbf{S}_{i_1, i_2, \dots, i_{p-2}}$ with respect to $\mathbf{X}$. These, combined with the fact that

$$D_{np}(0) = \min_{\mathbf{I}_{p-2} \in \mathcal{I}_{p-2}} D_{np}(\mathbf{S}_{\mathbf{I}_{p-2}}) - \frac{p-2}{n}$$

obtained in Proposition 1, motivate us to consider the following *iterative* procedure.

For convenience, in the sequel we assume that $\mathbf{X}^n \cap \{0\} = \emptyset$. Otherwise, using similar adjusted steps as Step 1-(a) and Step 2 in Algorithm 2 can remedy the final results.

Algorithm 3.

Input: $\mathbf{X}^n = \{\mathbf{x}_1, \dots, \mathbf{x}_n\} \in \mathcal{R}^p$, $3 \leq p < n < \infty$, $\mathbf{X}^n$ in general position.

Step 1. Set $d_0 = 1$, $Q = \emptyset$, $\mathcal{N} = \emptyset$.

Step 2. Compute $u_j = \mathbf{x}_j / \|\mathbf{x}_j\|$, $j = 1, \dots, n$. Set $\mathbf{U}_0 = \{u_1, -u_1, \dots, u_n, -u_n\}$, do:

(i) find $u_{\min} \in \mathbf{U}_0$ such that

$$u_{\min} = \arg \min_{u \in \mathbf{U}_0} P_n(u^\top X \leq 0),$$

(ii) compute $(i_{1,0}, i_{2,0}, \dots, i_{k,0}, i_{k+1,0}, \dots, i_{n,0})$ such that

$$u_{\min} \mathbf{x}_{i_{1,0}} \leq u_{\min} \mathbf{x}_{i_{2,0}} \leq \dots \leq u_{\min} \mathbf{x}_{i_{k,0}} < 0 \leq u_{\min} \mathbf{x}_{i_{k+1,0}} \leq \dots \leq u_{\min} \mathbf{x}_{i_{n,0}},$$

(iii) determine an $\mathbf{x}_{i_{p-1,0}}$ such that $D_{np}(\mathbf{S}_{i_{k+1,0}, i_{k+2,0}, \dots, i_{k+p-2,0}}) = P_n(u_0^\top X \leq 0)$ based on Algorithm 2, where u_0 is normal to the hyperplane passing through $\{0, \mathbf{x}_{i_{k+1,0}}, \mathbf{x}_{i_{k+2,0}}, \dots, \mathbf{x}_{i_{k+p-2,0}}, \mathbf{x}_{i_{p-1,0}}\}$, set $\mathbf{I}_{p-1} = \{i_{k+1,0}, \dots, i_{k+p-2,0}, i_{p-1,0}\}$, and store the depth $d_0 = D_{np}(\mathbf{S}_{i_{k+1,0}, i_{k+2,0}, \dots, i_{k+p-2,0}})$.

Step 3. (a) Push $\mathbf{I}_{p-1}$ into both $\mathcal{Q}$ and $\mathcal{N}$, (b) store the corresponding depth value $d_0 = D_{np}(\mathbf{S}_{\mathbf{I}_{p-1}})$.

Step 4. If $\mathcal{N}$ is non-empty, pop a combination $\mathbf{I}_{\text{cur}} = \{i_1, i_2, \dots, i_{p-1}\}$ from $\mathcal{N}$. For each $\{j_1, j_2, \dots, j_{p-2}\} \subset \mathbf{I}_{\text{cur}}$, do:

- (a) compute $D_{np}(\mathbf{S}_{j_1, j_2, \dots, j_{p-2}})$ by Algorithm 2,
- (b) store in $\mathcal{T}$ all $i_{p-1} \in \{1, 2, \dots, n\} \setminus \{j_1, j_2, \dots, j_{p-2}\}$ such that $\{0, \mathbf{x}_{j_1}, \mathbf{x}_{j_2}, \dots, \mathbf{x}_{j_{p-2}}, \mathbf{x}_{i_{p-1}}\}$ determine a u_0 satisfying $P_n(u_0^\top X \leq 0) - 1/n = D_{np}(\mathbf{S}_{j_1, j_2, \dots, j_{p-2}})$,
- (c) for each $i_{p-1} \in \mathcal{T}$, do: if $\mathbf{I}_{\text{new}} = \{j_1, j_2, \dots, j_{p-2}, i_{p-1}\}$ does not exist in $\mathcal{Q}$, store $D_{np}(\mathbf{S}_{j_1, j_2, \dots, j_{p-2}})$ and push $\mathbf{I}_{\text{new}}$ into both $\mathcal{Q}$ and $\mathcal{N}$,
- (d) if $d_0 > D_{np}(\mathbf{S}_{j_1, j_2, \dots, j_{p-2}}) - (p-2)/n$, set $d_0 = D_{np}(\mathbf{S}_{j_1, j_2, \dots, j_{p-2}}) - (p-2)/n$ and goto Step 5.

Step 5. Delete in both $\mathcal{Q}$ and $\mathcal{N}$ all combinations such that have Tukey depth values larger than d_0 .

Step 6. If $\mathcal{N}$ is not empty, then goto Step 4, else goto the next step.

Step 7. Obtain all data points, say $\mathbf{X}^s := \{\mathbf{x}_{k_l}\}_{l=1}^s$, lying on and cut off by the hyperplanes determined by $\{0\}$ and $\mathbf{I}_{p-1} \in \mathcal{Q}$. Similar to Algorithm 1, for each $\mathbf{I}_{p-2} := \{i_1, \dots, i_{p-2}\} \in \widetilde{\mathcal{I}}_{p-2} = \{\{j_1, j_2, \dots, j_{p-2}\} \mid j_1, j_2, \dots, j_{p-2} \text{ distinct, and } j_1, j_2, \dots, j_{p-2} \in \{k_1, k_2, \dots, k_s\}\}$, do:

- (a) compute $D_{np}(\mathbf{S}_{i_1, i_2, \dots, i_{p-2}})$,
- (b) if $d_0 > D_{np}(\mathbf{S}_{i_1, i_2, \dots, i_{p-2}}) - (p-2)/n$, set $d_0 = D_{np}(\mathbf{S}_{i_1, i_2, \dots, i_{p-2}}) - (p-2)/n$.

Step 8. Return d_0 .

Output: $D_{np}(0) = d_0$.

In Algorithm 3, *Step 2* are used mainly to computing an approximate Tukey depth d_0 and an initial combination $\mathbf{I}_{p-1}$. The direction vectors considered in *Step 2* are useful in reducing the computational burden if $D_{np}(0)$ is small. *Steps 4-7* are the key steps of Algorithm 3. Since Proposition 3 guarantees that the Tukey depth of each $\mathbf{S}_{j_1, j_2, \dots, j_{p-2}}$ considered in *Step 4* is no larger than that of $\mathbf{S}_{i_1, i_2, \dots, i_{p-2}}$, a great proportion of *critical direction vectors* would be *adaptively* eliminated from the computation. Furthermore, based on the former iteration, since we timely update in *Step 5* both Q and N by deleting the combinations $\{\mathbf{x}_{i_1}, \mathbf{x}_{i_2}, \dots, \mathbf{x}_{i_{p-2}}\}$ lie more inside with respect to the observations, Algorithm 3 requires quite minimum memory.

By design, in *Step 7*, all hyperplanes determined by $\{0\}$ and $\mathbf{I}_{p-1} \in Q$ form an envelope of the Tukey depth region of the value $D_{np}(\mathbf{S}_{\mathbf{I}_{p-1}}) (\geq D_{np}(0))$. They divide the observations into two parts. We use the part containing less observations to compute the Tukey depth with complexity less than $O\left(\frac{1}{2^{p-2}} n^{p-1} \log(n)\right)$.

5 Performances

In this section, we will conduct a few data examples to investigate the performance of the proposed algorithms. All of these results are obtained on a HP Pavilion dv7 Notebook PC with Intel(R) Core(TM) i7-2670QM CPU @ 2.20GHz, RAM 6.00GB, Windows 7 Home Premium and Matlab 7.8.

5.1 Illustrations

In this subsection, we are interested to illustrate the performances of the proposed algorithms in terms of both computation time and accuracy based on the real data. For the sake of comparison, we also report the results obtained by the combinatorial algorithm developed by Mozharovskiy (2014), and the naive algorithm proposed by Liu and Zuo (2014a). For convenience, in the sequel we denote the refine combinatorial algorithm as *RCom*, the adaptive iterative algorithm as *ADIA*, and the algorithms of Mozharovskiy (2014) and Liu and Zuo (2014a) as *DM14* and *LZ14*, respectively.

Two data sets are considered in the following. The first data set is taken from Härdle and Simar (2007), and has been investigated by Liu and Zuo (2014a) as an illustration. It consists of 64 samples as a part of an evolution of the vocabulary of children obtained from a cohort of pupils from the eighth through 11th grade levels. The second data set is a part of the daily simple returns of IBM stock from 1970 January 01 to 2008 December 25. It currently can be downloaded from his teaching page: <http://faculty.chicagobooth.edu/ruey.tsay/teaching>

The original data set consists of 755 observations; see [Tsay \(2010\)](#) for more details. Remarkably, our goal here is not to perform a thorough analysis for data, but rather to show how the algorithms work in practice.

Table 1: Computation time (in seconds).

p	Data	n	Computation time					
			$RCom$	$ADIA_{\min}$	$ADIA_{\text{mean}}$	$ADIA_{\max}$	$DM14$	$LZ14$
3	Voc	64	0.0145	0.0066	0.0117	0.0380	0.1108	2.7740
	IBM	200	0.0908	0.0146	0.0387	0.1287	1.1892	31.9022
	IBM	500	0.4925	0.0283	0.1472	0.5609	8.2659	285.4298
4	Voc	64	0.4614	0.0335	0.1410	0.3731	2.9491	89.8435
	IBM	200	8.8961	0.0620	0.4624	3.3255	86.4703	3476.2415
	IBM	500	120.0689	0.1411	2.4635	21.9474	1721.5922	--

Both data sets are 4-dimensional. Using the proposed algorithms both obtain the *exact* Tukey depth, which coincide with those computed by $DM14$ and $LZ2014$. Table 1 reports the computation time (in seconds) for calculating the Tukey depth of a single observation. Here Voc denotes the vocabulary data, and IBM stands for the IBM stock data. $p = 3$ means we only use the first three columns of the data set, and $n = 200$ the first 200 rows. The sign ‘--’ in Table 1 means this depth value is not computable in 8 hours. Since all $Rcom$, $DM14$ and $LZ14$ have to fully address $\binom{n}{p-1}$ critical direction vectors for every observation, the time for calculating each observation is almost the same. We only list the average computation time. Whereas the computation time consumed by $ADIA$ depends on the Tukey depth of the point being computing for a given data set, and therefore we report additionally its minimum, mean and maximum computation time (under the titles $ADIA_{\min}$, $ADIA_{\text{mean}}$ and $ADIA_{\max}$, respectively). The smaller the depth value of the observation is, the less the computation time the algorithm $ADIA$ tends to consume; see Figure 2 for more details.

Table 1 indicates that both the proposed algorithms run much faster than the existing algorithms. Among them, the implementation of $ADIA$ tends to run most the fastest when n and/or p are large. It requires no more than 3 seconds (in average) to obtain the depth of a single point in all illustrations here.

It is worth mentioning that the algorithm of [Mozharovskiy \(2014\)](#) is implemented here by us in Matlab for convenience of comparison. It appears to be slower than what have been reported in [Mozharovskiy \(2014\)](#). This is possible, because their computations are based on C++, which usually runs faster than Matlab, especially when there are a great number of iterations involved.




Figure 2: Shown are the Tukey depth values (sorted ascending) and the corresponding computation times (in seconds) of the observations of Voc and IBM consumed by ADIA.

5.2 Speed comparisons

In the following, we further compare the speeds of the proposed algorithms with that of *DM14* based on the simulated data. The data are generated from the 3, 4, 5, 6-dimensional standard normal distributions with sample size $n = 40, 80, 160, \dots, 2560$. For each combination of $p \in \{3, 4, 5, 6\}$ and $n \in \{40, 80, 160, \dots, 2560\}$, we compute repeatedly 10 times the Tukey depth values of $z = \alpha \mathbf{1}_p$ with $\alpha = 0, 0.4, 0.8, 1.2$, where $\mathbf{1}_p$ denotes the p -dimensional vector of ones. We report the average computation time of *RCom* and *DM14* in Table 2, and that of *ADIA* in Table 3, respectively. Since as pointed above, the

computation time of both *RCom* and *DM14* do not depend on the Tukey depth of $z = 0$ being computing, we report only the average computation time corresponding to $\alpha = 0$ here. Tables 2-3 indicate that both the proposed algorithms run much faster than that of *DM14*.

Table 2: Average computation times (in seconds) of *RCom* and *DM14*.

p	Method	n						
		40	80	160	320	640	1280	2560
3	$RCom$	0.0268	0.0481	0.0899	0.2531	0.8073	4.6438	11.7662
	$DM14$	0.1466	0.8860	5.1369	35.0182	255.9098	2072.8847	16230.1634
4	$RCom$	0.1876	0.8045	5.0550	43.3812	248.5797	1878.8783	14681.9140
	$DM14$	0.6377	5.7872	45.1497	393.0911	3612.5004	--	--
5	$RCom$	1.9517	21.1227	262.3251	3864.4624	--	--	--
	$DM14$	5.8554	104.6715	1847.4277	--	--	--	--
6	$RCom$	18.4515	446.0153	14695.1294	--	--	--	--
	$DM14$	48.6214	1794.7799	--	--	--	--	--

Table 3: Average computation times (in seconds) of *ADIA*.

p	α	n						
		40	80	160	320	640	1280	2560
3	0.0	0.0193	0.0634	0.0503	0.1297	0.4372	1.8128	9.1473
	0.4	0.0167	0.0470	0.0668	0.0968	0.3418	1.3816	5.3099
	0.8	0.0065	0.0131	0.0180	0.0460	0.3248	0.4342	5.4500
	1.2	0.0067	0.0136	0.0186	0.0390	0.1220	0.3514	1.3141
4	0.0	0.3416	0.3211	1.7991	6.5374	60.6911	408.8870	5570.4364
	0.4	0.0607	0.1704	0.5131	1.7255	11.9404	105.2564	740.5502
	0.8	0.1070	0.1172	0.1140	0.5171	1.7526	10.1355	72.9394
	1.2	0.1373	0.1465	0.0538	0.2278	0.9409	2.2443	5.0238
5	0.0	0.9466	6.9212	68.0360	404.5147	13177.5227	--	--
	0.4	0.4048	1.0052	8.3791	43.6717	609.6461	11022.4774	--
	0.8	1.2751	1.1352	1.6600	5.3665	15.8776	105.1733	1749.6048
	1.2	0.1923	0.1001	0.3339	0.3162	1.5744	47.5422	91.6245
6	0.0	12.3953	95.3378	3883.9385	--	--	--	--
	0.4	4.3607	18.3686	173.8880	1504.1087	--	--	--
	0.8	0.1638	0.2295	62.8009	158.7744	1777.3669	--	--
	1.2	1.1149	3.0121	1.4400	4.9680	19.3288	30.2152	75.0221

6 Concluding discussions

In this paper, we considered the computing issue of the Tukey depth. We found that it is useful to extend the conventional definition of the Tukey depth for a single point

into the version for a subspace. Based on this extended definition, we suggested two new algorithms for fast computing the Tukey depth. That is, a refined combinatorial algorithm (*Rcom*) and an adaptive iterative algorithm (*ADIA*).

Among them, *Rcom* is quite naive. It is in fact a refined version of the combinatorial algorithm (*DM14*) suggested by Mozharovskiy (2014). The complexity of *Rcom* is exactly $O(n^{p-1} \log(n))$, lower than $O(n^p)$ of *DM14*. Note that when computing the Tukey depth, we are in fact searching for the infimum of $P_n(u^\top X \leq u^\top z)$ with respect to u . Hence, many *critical direction vectors* addressed by *Rcom* are *redundant*, and could be eliminated. *ADIA* further takes into account this fact through using the *quasiconcave* of the Tukey depth function. Therefore, *ADIA* can be implemented more efficiently than *Rcom* with complexity less than $O(n^{p-1} \log(n)/2^{p-2})$.

From both *Rcom* and *ADIA*, it is easy to see that it is a very key step to compute the bivariate Tukey depth, because it would be repeatedly computed considerable times when n and/or p are large. In the literature, it seems the best procedure of computing the bivariate Tukey depth of a single point has only the complexity $\log^2(n)$ (Miller et al., 2003). Hence, the proposed algorithms may be further improved thought *simply replacing Algorithm 2 with the more efficient procedure*.

As mentioned in Mozharovskiy (2014), there are many depth notions, similar to the Tukey depth, is of both projection and *quasiconcave* properties, such as the projection depth (Zuo, 2003) and the zonoid depth (Koshevoy and Mosler, 1997). Efficient algorithms exist only for bivariate data for these depth notions; see for example Liu and Zuo (2014b). Therefore, how to utilize similarly the *quasiconcave* of these depth notions to reduce the computational burden in higher dimensions is still worthy of further consideration.

Acknowledgments

The author thanks Prof. Mosler, K. and Dr. Mozharovskiy, P. for their valuable discussions during the preparation of this manuscript. This research is partly supported by the National Natural Science Foundation of China (No.11461029), the Natural Science Foundation of Jiangxi Province (No.20142BAB211014), and the Youth Science Fund Project of Jiangxi provincial education department (No. GJJ14350).

References

Edelsbrunner, H., 1987. *Algorithms in Combinatorial Geometry*. Springer, Heidelberg.

- Hallin, M., Paindaveine, D., Šiman, M., 2010. Multivariate quantiles and multiple-output regression quantiles: From L_1 optimization to halfspace depth. *Ann. Statist.* 38, 635-669.
- Härdle, W., and Simar, L. (2007). *Applied Multivariate Statistical Analysis*. Springer, Heidelberg.
- Johnson, T., Kwok, I. and Ng, R. (1998). Fast computation of 2-dimensional depth contours. In: Agrawal, R., Stolorz, P. (eds.), *Proceedings of the Fourth International Conference on Knowledge Discovery and Data Mining*, AAAI Press, New York, 224-228.
- Kong, L., Mizera, I., 2012. Quantile tomography: Using quantiles with multivariate data. *Statist. Sinica*, 22, 1589-1610.
- Kong, L., Zuo, Y., 2010. Smooth depth contours characterize the underlying distribution. *J. Multivariate Anal.*, 101, 2222-2226.
- Koshevoy, H., Mosler, K., 1997. Zonoid trimming for multivariate distributions. *Ann. Statist.* 25, 1998-2017.
- Li, J., Cuesta-Albertos, J.A., Liu, R.Y., 2012. DD-classifier: nonparametric classification procedure based on DD-plot. *J. Amer. Statist. Assoc.* 107(498), 737-753.
- Liu, X., Zuo, Y., 2014a. Computing halfspace depth and regression depth. *Communications in Statistics-Simulation and Computation*, 43, 969-985.
- Liu, X.H., Zuo, Y.J., 2014b. Computing projection depth and its associated estimators. *Statistics and Computing*, 24(1), 51-63.
- Miller, K., Ramaswami, S., Rousseeuw, P., Sellars, J. A., Souvaine, D., Streinu, I., Struyf, A. 2003. Efficient computation of location depth contours by methods of computational geometry. *Statistics and Computing*, 13(2), 153-162.
- Mosler, K. 2013. Depth statistics. In *Robustness and Complex Data Structures* (pp. 17-34). Springer Berlin Heidelberg.
- Mosler, K., Lange, T., Bazovkin, P., 2009. Computing zonoid trimmed regions of dimension $d > 2$. *Comput. Statist. Data Anal.* 53, 2500-2510.
- Mozharovskiy, P., 2014. Contributions to depth-based classification and computation of halfspace depth, PhD thesis, http://www.wisostat.uni-koeln.de/fileadmin/sites/statistik/pdf_publikationen/PhD-thesis_Mozharovskiy.
- Rousseeuw, P.J., Ruts, I., 1996. Algorithm AS 307: Bivariate location depth. *J. Appl. Statist.* 45, 516-526.
- Rousseeuw, P.J., Struyf, A., 1998. Computing location depth and regression depth in higher dimensions. *Statist. Comput.* 8, 193-203.
- Tsay, R.S., 2010. *Analysis of Financial Time Series*, Third Edition. John Wiley & Sons.
- Tukey, J.W., 1975. Mathematics and the picturing of data. In *Proceedings of the International Congress of Mathematicians*, 523-531. Cana. Math. Congress, Montreal.
- Yeh, A., Singh, K., 1997. Balanced confidence regions based on Tukey's depth and the bootstrap. *J. Roy. Statist. Soc. Ser. B* 59, 639-652.
- Zuo, Y.J., 2003. Projection based depth functions and associated medians. *Ann. Statist.* 31, 1460-1490.
- Zuo, Y.J., Serfling, R., 2000. General notions of statistical depth function. *Ann. Statist.* 28, 461-482.